



Immune Response in Head Kidney of Rainbow Trout Fry Following Stress and Infection with *Flavobacterium Psychrophilum*

Henriksen, Maya Maria Mihályi; Kania, P. W.; Madsen, Lone; Buchmann, K.; Dalsgaard, Inger

Publication date:
2013

[Link back to DTU Orbit](#)

Citation (APA):

Henriksen, M. M. M., Kania, P. W., Madsen, L., Buchmann, K., & Dalsgaard, I. (2013). *Immune Response in Head Kidney of Rainbow Trout Fry Following Stress and Infection with Flavobacterium Psychrophilum*. Abstract from 16th International Conference on Diseases of Fish and Shellfish, Tampere, Finland.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

IMMUNE RESPONSE IN HEAD KIDNEY OF RAINBOW TROUT FRY FOLLOWING STRESS AND INFECTION WITH *FLAVOBACTERIUM PSYCHROPHILUM*.

M. M. M. Henriksen^{*1}, P. W. Kania², L. Madsen¹, K. Buchmann² and I. Dalsgaard¹

¹*National Veterinary Institute, Technical University of Denmark, Copenhagen, Denmark*

²*Department of Veterinary Disease Biology, University of Copenhagen, Copenhagen, Denmark*

The bacterial fish pathogen *Flavobacterium psychrophilum*, the cause of Rainbow Trout Fry Syndrome (RTFS), results in significant mortality in farmed rainbow trout; unless it is treated with antibiotics. Presently no commercial vaccine exists. More knowledge is required to elucidate the immune response in rainbow trout against *F. psychrophilum* in order to create preventive measures against RTFS. A limited number of studies have been carried out so far and have relied on samples from either naturally infected or injection-challenged fish. The use of naturally infected fish introduces many possible sources of error. Injection is a suboptimal approach for investigations regarding the immune response, since mucosal surfaces are bypassed. *F. psychrophilum* has a limited ability to cause disease in experimental bath challenges without applying a stressor. Recently, a bath model utilized H₂O₂ before pathogen exposure to elevate mortality. The model was used to examine the immune response to infection in rainbow trout fry (≈ 1 g); both with and without preceding H₂O₂ treatment. Samples from the head kidney were taken before pathogen exposure and 4 hours, 48 hours, 125 hours and 192 hours after exposure. The regulation of several immune relevant genes was examined and the relative bacterial load was assessed. Although it is not determined how H₂O₂ increases mortality, it is assumed to be due to stress. Exposure to H₂O₂ prior to infection altered the regulation of several genes, and several correlations between pathogen load and gene expression were observed.